**Few-Body Systems manuscript No.**
(will be inserted by the editor)

**Loredana Bellantuono**

# On $J^{PC} = 0^{--}$ exotic glueball

arXiv:1602.01349v1 [hep-ph] 3 Feb 2016

Received: date / Accepted: date

**Abstract** The mass spectrum of the gluonium with $J^{PC} = 0^{--}$ is examined in three bottom-up AdS/QCD models. The results are used to identify several production and decay modes useful for searching this state. Moreover, the properties of such glueball in a hot and dense quark medium are discussed.

**Keywords** QCD Phenomenology · Glueball and nonstandard multi-quark/gluon states

**PACS** 11.25.Tq · 11.10.Kk · 12.39.Mk

## 1 Introduction

Glueballs, bound states of gluons arising from the non-Abelian nature of strong interactions, are an important testbed for non perturbative aspects of QCD. The main obstacle in their search is the mixing with quarkonia $(\bar{q}q)$ with the same quantum numbers. A promising strategy for their identification is to focus on exotic states whose $J^{PC}$ quantum numbers are unaccessible to quark-antiquark configurations. This is the case of several gluonia with negative $C$-parity, composed of an odd number of constituent gluons ("oddballs"), for which little theoretical information is available. In particular, for $J^{PC} = 0^{--}$, the mass predictions for the lightest state span the range from $m_{0^{--}} = 2.79$ GeV in the flux-tube model [1], to $m_{0^{--}} \approx 5.166$ GeV in lattice QCD simulations [2]. Two stable $0^{--}$ oddballs, with masses $m_{0^{--}} = 3.81 \pm 0.12$ GeV and $m_{0^{--}} = 4.33 \pm 0.13$ GeV, have been predicted by QCD sum rules [3].

The mass spectrum of the $J^{PC} = 0^{--}$ oddball can be computed in a framework inspired by the AdS/CFT correspondence. This duality conjecture relates a strongly coupled gauge theory in a four dimensional $(4D)$ Minkowski space to a semiclassical gravity theory defined in a five dimensional $(5D)$ anti-de Sitter (AdS) geometry times a $5D$ sphere [4]. In Poincaré coordinates, the line element

$$ds^2 = \frac{R^2}{z^2}(dx_0^2 - d\mathbf{x}^2 - dz^2) \qquad z > 0\,, \tag{1}$$

with $z$ the fifth holographic coordinate, describes the AdS bulk metric. The original formulation of gauge/gravity duality required the $4D$ theory to be conformal invariant. Holographic bottom-up models, constructed to reproduce QCD properties, break such a symmetry by introducing an infrared energy scale in the bulk; the $J^{PC} = 0^{--}$ oddball mass spectrum can then be determined either in vacuum or in a quark bath at finite temperature and density. Such an approach complements the top-down methods applied to analyze, e.g., the scalar $0^{++}$ gluonium [5; 6].

L. Bellantuono
Dipartimento di Fisica, Università di Bari, and INFN, Sezione di Bari, via Orabona 4, I-70126 Bari, Italy
E-mail: loredana.bellantuono@ba.infn.it

## 2 $J^{PC}=0^{--}$ mass spectrum in three holographic models of QCD

The holographic correspondence is based on a dictionary, according to which a local gauge-invariant operator in the $4D$ field theory is dual to a field in $5D$ AdS [7; 8]. In QCD, an interpolating current representing the glueball with quantum numbers $J^{PC}=0^{--}$ can be written in terms of the gluon field strengths $G^a_{\mu\nu}(x)$ and $\widetilde{G}^a_{\mu\nu}(x)=\frac{1}{2}\varepsilon_{\mu\nu\rho\sigma}G^a_{\rho\sigma}(x)$, namely

$$J_0(x)=g_s^3 d_{abc}[\eta^t_{\alpha\beta}\widetilde{G}^a_{\mu\nu}(x)][\partial_\alpha\partial_\beta G^b_{\nu\rho}(x)][G^c_{\rho\mu}(x)]\,, \tag{2}$$

with $a,b,c$ color indices, $d_{abc}$ the symmetric tensor defining the anticommutator of $SU(3)_c$ generators, and $g_s$ the strong coupling constant. The transverse $\eta^t_{\alpha\beta}$ metric is defined as $\eta^t_{\alpha\beta}=\eta_{\alpha\beta}-\frac{\partial_\alpha\partial_\beta}{\partial^2}$, with $\alpha,\beta$ (as well as $\mu,\nu$) $4D$ Lorentz indices, and $\eta_{\alpha\beta}$ the Minkowski metric tensor [3; 9]. The operator (2) has conformal dimension $\Delta=8$, and its holographic dual field, $O_0(x,z)$, has mass obtained by the relation $M_5^2R^2=\Delta(\Delta-4)$ [7; 8]. In the following, the $AdS_5$ radius is set to $R=1$. The $5D$ action for $O_0(x,z)$ can be written as

$$S=\frac{1}{k}\int d^5x\sqrt{g}a(z)\left[g^{MN}\partial_M O_0\,\partial_N O_0-M_5^2O_0^2\right]\,, \tag{3}$$

where $g_{MN}$ is the bulk metric, $g=|det(g_{MN})|$ and $k$ is a parameter making the action dimensionless. To account for confinement in QCD, conformal invariance must be broken in the action (3). Three different models including such effect, either by a proper choice of the function $a(z)$ or by introducing a dynamical field in the metric, are discussed in the following. The mass spectrum can be determined solving the Euler-Lagrange equations for the field $O_0(x,z)$.

*Hard-wall model.* A simple way of modeling confinement in the holographic setup is by considering a slice of the $AdS_5$ space, with a sharp cutoff at a finite distance $z_m$ along the fifth dimension [10]. The metric $g_{MN}$ is given by (1), and the condition $z\leq z_m$ is implemented in the action through $a(z)=\Theta(z_m-z)$, with $\Theta$ the Heaviside function. The cutoff $z_m$ sets a mass scale. The choice $1/z_m=346$ MeV, obtained from analyses on axial and vector mesons [10], gives the results $m_0=2.80$ GeV and $m_1=4.14$ GeV for the lowest-lying and the first-excited $0^{--}$ oddball, respectively [11].

*Soft-wall model.* In this framework, the geometry is $AdS_5$ (1), and conformal invariance is smoothly broken by the function $a(z)=e^{-c^2z^2}$ which introduces a mass scale $c$ in the action [12]. The Regge-like mass spectrum is obtained [11]:

$$m_n^2=4c^2(n+4)\,. \tag{4}$$

Setting $c=m_\rho/2=388$ MeV from the $\rho$ meson mass computed in the model [12], one obtains $m_0=1.55$ GeV and $m_1=1.74$ GeV [11]. The mass spectrum (4) corresponds to the poles of the two-point correlation function of $J_0(x)$,

$$\Pi(p^2)=i\int d^4x\,e^{ipx}\langle 0|T[J_0(x)J_0^\dagger(0)]|0\rangle\,. \tag{5}$$

In the AdS/CFT dictionary, the QCD interpolating current $J_0(x)$ is interpreted as the source of the dual field $O_0(x,z)$. The $4D$ Fourier transforms of such operators, $\tilde{J}_0(p)$ and $\tilde{O}_0(p,z)$, are related by the bulk-to-boundary propagator $\tilde{K}(p,z)$ of the oddball field, through $\tilde{O}_0(p,z)=\tilde{K}(p,z)\tilde{J}_0(p)$. Holography prescribes the identification between the partition functions of the dual theories, and this allows to compute the two-point correlation function (5) as

$$\Pi\left(p^2\right)\approx\left.\frac{\delta^2 S_{os}}{\delta\tilde{J}_0\delta\tilde{J}_0}\right|_{\tilde{J}_0=0}\,, \tag{6}$$

with $S_{os}$ the $5D$ on-shell action. The poles of $\Pi\left(p^2\right)$ are given in (4).

**Table 1** Production and decay modes of the $J^{PC}=0^{--}$ oddball, for $m_{0^{--}}=2.8$ GeV.

| Radiative production | Hadronic production | Decay mode |
|---|---|---|
| $\chi_{c1}(3510)\to\gamma G(0^{--})$ | $X(3872)\to\omega G(0^{--})$ | $G(0^{--})\to\gamma f_1(1285)$ |
| $X(3872)\to\gamma G(0^{--})$ | $h_c(3525)\to\pi\pi\,(I=0)\,G(0^{--})$ | $G(0^{--})\to\omega f_1(1285)$ |
| $\chi_{c2}(3556)\to\gamma G(0^{--})$ | $\chi_{b1}(10255)\to(\omega,\phi,J/\Psi)\,G(0^{--})$ | $G(0^{--})\to\rho\, a_1(1260)\ (I=0)$ |
| $\chi_{c2}(3927)\to\gamma G(0^{--})$ | $\Upsilon(nS)\to(f_1(1285),\chi_{c1},X(3872))\,G(0^{--})$ | |
| $\chi_{b1}(9892)\to\gamma G(0^{--})$ | $h_b(9899)\to f_0(980)\,G(0^{--})$ | $G(0^{--})\to h_1(1270)\,f_0(980)$ |
| $\chi_{b1}(10255)\to\gamma G(0^{--})$ | $h_b(10260)\to f_0(980)\,G(0^{--})$ | $G(0^{--})\to\rho\,\pi\ (I=0)$ |
| $\chi_{b2}(9912)\to\gamma G(0^{--})$ | $h_b(9899)\to G(0^{++})\,G(0^{--})$ | $G(0^{--})\to K^*K\ (I=0)$ |
| $\chi_{b2}(10269)\to\gamma G(0^{--})$ | $h_b(10260)\to G(0^{++})\,G(0^{--})$ | $G(0^{--})\to(\eta,\eta')(\omega,\phi)$ |

*Einstein-dilaton model.* A third possibility is to consider a class of dynamical bottom-up models reproducing confinement through a distorsion of the bulk geometry [13; 14]:

$$ds^2_{(ED)}=\frac{e^{2A_s(z)-\frac{4}{3}\Phi(z)}}{z^2}\left[dx_0^2-d\mathbf{x}^2-dz^2\right]. \tag{7}$$

$\Phi(z)$, a dilaton field, couples to the graviton, and its profile is determined solving the Einstein equations deduced from this metric. The function $A_s(z)$ can be chosen as $A_s(z)=\delta^2z^2$, with the mass scale fixed to $\delta=0.43$ GeV [13]. Once the profile of $\Phi(z)$ is obtained, the two lightest oddball states turn out to have mass $m_0=2.82$ GeV and $m_1=4.07$ GeV [11].

With these results, several production and decay modes can be identified, useful for the search of the $0^{--}$ gluonium. They are reported in Table 1 for the specific case $m_{0^{--}}=2.8$ GeV [11].

## 3 Oddball in thermalized and dense medium

The AdS/QCD duality allows to examine the properties of a $J^{PC}=0^{--}$ oddball in a quark thermal bath, at finite temperature $T$ and chemical potential $\mu$. Stability of the gluon configuration against thermal and density fluctuations can be investigated in comparison with other quark and gluon bound states. In-medium effects can be incorporated in the holographic description using the action (3), with an appropriate bulk geometry. I explicitly discuss the soft-wall case. The $5D$ line element

$$ds^2=\frac{1}{z^2}\left(f(z)dx_0^2-d\mathbf{x}^2-\frac{dz^2}{f(z)}\right) \tag{8}$$

is characterized by a metric with $f(z)$ which can be related to temperature and density of the boundary theory. The $5D$ Reissner-Nordström $AdS$ geometry (AdS/RN), with

$$f(z)=1-\left(\frac{1}{z_h^4}+q^2z_h^2\right)z^4+q^2z^6, \tag{9}$$

can be used as the holograpic dual of a thermalized and dense medium in the deconfined phase [15; 16]. This charged black hole bulk metric has an outer horizon $z=z_h$ and charge $q$ which can be related to the temperature $T$ and chemical potential $\mu$ of the dual boundary theory. Defining $Q=qz_h^3$ and imposing the condition $0\le Q\le\sqrt{2}$, the black-hole temperature is

$$T=\frac{1}{4\pi}\left|\frac{df}{dz}\right|_{z=z_h}=\frac{1}{\pi z_h}\left(1-\frac{Q^2}{2}\right). \tag{10}$$

In the QCD generating functional, $\mu$ multiplies the quark number operator $O_q(x)=q^\dagger(x)q(x)$. Thus, it can be interpreted, according to gauge/gravity correspondence, as the source of a bulk field dual to $O_q(x)$, the time component of a $U(1)$ gauge field $A_M(x,z)$. By rotational invariance, the spatial components $A_i$ (with




**Fig. 1** Spectral function $\rho\left(\omega^2\right)/\left(\omega^2\right)^6$ computed in the soft-wall model with AdS/RN metric, at fixed $\mu$ (left) or $T$ (right). The dimensionful quantities are in units of $c$, and the constant $10^{-8}k/\left(2z_h^8\right)$ has been factorized out in the spectral function [11].




**Fig. 2** Squared mass $m^2$ and width $\Gamma$ of the lightest $J^{PC}=0^{--}$ glueball, obtained from the soft-wall model with AdS/RN metric, in a range of temperature $T$ and chemical potential $\mu$, in units of $c$ [11].

$i=1,2,3,z$) vanish, while $A_0\approx\mu-\kappa qz^2$ for small $z$. The condition $A_0(z_h)=0$ provides a linear relation between $\mu$ and $q$:

$$\mu=\kappa\frac{Q}{z_h}\,, \tag{11}$$

with $\kappa$ a dimensionless parameter that will be fixed to 1, a choice determining the quark chemical potential $\mu$ up to a numerical factor. Important information on the QCD bound states in a quark thermal bath can be inferred from the spectral function $\rho\left(\omega^2\right)$, the imaginary part of the retarded Green function. The spectral function of the $0^{--}$ oddball is represented in vacuum by an infinite number of delta functions centered at the eigenvalues of the mass spectrum (4). Fig. 1 shows the changes of the in-medium $\rho\left(\omega^2\right)$ as temperature and chemical potential are switched on. At any finite and fixed value of $\mu$, the peaks of the spectral function broaden and move towards lower values of $\omega^2$ when $T$ increases. An analogous behaviour is observed by keeping the temperature fixed, and increasing the chemical potential. The $(T,\mu)$-dependence of the lightest oddball's squared mass and width is shown in Fig. 2. The values of the temperature and chemical potential at which the lowest-energy peak becomes indistinguishable in the profile of the spectral function, are smaller than those obtained for light vector [17; 18], scalar $\bar{q}q$ mesons [19], the lightest scalar glueball [20], and hybrid mesons [21]. Hence, $0^{--}$ oddballs are found to suffer of larger in-medium instabilities with respect to other hadrons.

For points in the $T-\mu$ plane below the QCD deconfinement transition, the state described by AdS/RN metric is metastable [16]. The thermal-charged AdS metric (tcAdS), with line element (8) and deformation

$$f(z)=1+q^2z^6\,. \tag{12}$$

is proposed as a dual of the confined phase of QCD at small temperature and finite chemical potential [16; 22]. This geometry is related to the AdS/RN one by the Hawking-Page transition, which describes deconfinement in the holographic framework. The chemical potential, related to the finite density of the hadronic medium,




**Fig. 3** Squared mass $m^2$ as a function of $\mu$ for the two lightest $J^{PC} = 0^{--}$ oddballs, in the tcAdS soft-wall model.

grows linearly with $q$, while the temperature is implemented through a periodicity in the Euclidean time coordinate $\tau = ix_0$ [22]. The results, depicted in Fig. 3, reveal that the masses of the two lightest oddballs increase with $\mu$, a different behaviour with respect to what is found in the deconfined phase [11].

## 4 Conclusions

In AdS/QCD, the mass of the lowest-lying $0^{--}$ oddball is found to be lighter than as computed by other approaches. The effect of deconfined quark matter can be modeled by the AdS/RN geometry: oddballs are found to suffer from larger thermal and density instabilities with respect to other hadrons. On the other hand, the holographic description of a confined medium, obtained using the tcAdS metric, gives a mass which increases with the chemical potential.

**Acknowledgements** I thank P. Colangelo and F. Giannuzzi for collaboration, and F. De Fazio and S. Nicotri for discussions.

## References

1. Isgur, N., Paton, J.E.: A Flux Tube Model for Hadrons in QCD. Phys. Rev. D **31**, 2910 (1985)
2. Gregory, E. et al.: Towards the glueball spectrum from unquenched lattice QCD. JHEP **1210**, 170 (2012)
3. Qiao, C.-F., Tang, L.: Finding the $0^{--}$ Glueball. Phys. Rev. Lett. **113**, 221601 (2014)
4. Maldacena, J.M.: The Large N limit of superconformal field theories and supergravity. Int. J. Theor. Phys. **38**, 1113 (1999)
5. Csaki, C., Ooguri, H., Oz, Y., Terning, J.: Glueball mass spectrum from supergravity. JHEP **9901**, 017 (1999)
6. Brunner, F., Rebhan, A.: Nonchiral enhancement of scalar glueball decay in the Witten-Sakai-Sugimoto model. Phys. Rev. Lett. **115**, 131601 (2015)
7. Witten, E.: Anti-de Sitter space and holography. Adv. Theor. Math. Phys. **2**, 253 (1998)
8. Gubser, S., Klebanov, I.R, Polyakov, A.M.: Gauge theory correlators from noncritical string theory. Phys. Lett. B **428**, 105 (1998)
9. Qiao, C.-F., Tang, L.: Mass Spectra of $0^{+-}$, $1^{-+}$, and $2^{+-}$ Exotic Glueballs. arXiv:1509.00305 [hep-ph] (2015)
10. Erlich, J. et al.: QCD and a holographic model of hadrons. Phys. Rev. Lett. **95**, 261602 (2005)
11. Bellantuono, L., Colangelo, P., Giannuzzi, F.: Holographic Oddballs. JHEP **1510**, 137 (2015)
12. Karch, A., Katz, E., Son, D.T., Stephanov, M.A.: Linear confinement and AdS/QCD. Phys. Rev. D **74**, 015005 (2006)
13. Li, D., He, S., Huang, M., Yan, Q.-S.: Thermodynamics of deformed $AdS_5$ model with a positive/negative quadratic correction in graviton-dilaton system. JHEP **1109**, 041 (2011)
14. Chen, Y., Huang, M.: Two-gluon and trigluon glueballs from dynamical holography QCD. arXiv:1511.07018 [hep-ph] (2015)
15. Colangelo, P., Giannuzzi, F., Nicotri, S.: In-medium hadronic spectral functions through the soft-wall holographic model of QCD. JHEP **1205**, 076 (2012)
16. Lee, B.-H., Park, C., Sin, S.-J.: A Dual Geometry of the Hadron in Dense Matter. JHEP **0907**, 087 (2009)
17. Fujita, M., Fukushima, K., Misumi, T., Murata, M.: Finite-temperature spectral function of the vector mesons in an AdS/QCD model. Phys. Rev. D **80**, 035001 (2009)
18. Giannuzzi, F.: Holographic vector mesons in a hot and dense medium. Nucl. Phys. Proc. Suppl. **234**, 309 (2013)
19. Colangelo, P., De Fazio, F., Giannuzzi, F., Jugeau, F. et al.: Light scalar mesons in the soft-wall model of AdS/QCD. Phys. Rev. D **78**, 055009 (2008)
20. Colangelo, P., De Fazio, F., Jugeau, F., Nicotri, S.: On the light glueball spectrum in a holographic description of QCD. Phys. Lett. B **652**, 73 (2007)
21. Bellantuono, L., Colangelo, P., Giannuzzi, F.: Exotic $J^{PC} = 1^{-+}$ mesons in a holographic model of QCD. Eur. Phys. J C **74**, 2830 (2014)
22. Park, C. et al.: The Soft Wall Model in the Hadronic Medium. Phys. Rev. D **84**, 046007 (2011)